

Nonstationary Aharonov-Bohm Effect of Current Correlation by Entangled Electrons in Quantum Spin Hall Systems

Wei Chen,^{1,2} Z. D. Wang,^{2,*} R. Shen,^{1,†} and D. Y. Xing¹

¹*National Laboratory of Solid State Microstructures and Department of Physics, Nanjing University, Nanjing 210093, China*

²*Department of Physics and Center of Theoretical and Computational Physics, The University of Hong Kong, Pokfulam Road, Hong Kong, China*

We investigate spin entangled electron pairs transporting separately in the helical edge channels of two quantum spin Hall systems. Both *dc* and *ac* Aharonov-Bohm (AB) effects of the crossed current correlation are studied. In the *dc* case, it is found that the visibility of the AB oscillation of differential noise is equal to entanglement concurrence, and a π phase difference indicates singlet and triplet entangled states. More remarkably, when an *ac* flux with frequency Ω is introduced, the differential noise exhibits nontrivial step structures as the voltage bias V increases, with the step edges being located at $eV = n\hbar\Omega$. The concurrence can be probed directly by measuring the heights of these steps. We also show that the relation between differential noise and concurrence is general even when spin dephasing exists.

PACS numbers: 03.65.Ud, 72.70.+m, 73.23.-b, 73.20.-r

As is known, entanglement reflects a kind of nonlocal correlation [1, 2]. The research on quantum entanglement has played an important role in quantum information and computation science [3]. Recently, the creation and detection of electronic entanglement in solid state systems have attracted much interest, for the large-scale implementation of quantum information and computation schemes [4]. Moreover, the crossed Andreev reflection in mesoscopic s-wave superconductor systems has already been confirmed [5], which is one of the main proposals for generation of spin entangled electrons in solids [6].

Though the idea to utilize entanglement in solid systems is elegant, a direct experimental evidence is still challenging. Several proposals for spin entanglement detection in solids have been put forward, including the Bell inequality tests [7, 8] and the shot noise in a beam splitter setup [9]. The former is based on local hidden variable theories [2] and the latter utilizes the relation between spin entanglement and antisymmetry of electronic wave functions [9]. Most recently, we have exploited another method [10], based on a completely different mechanism, so called quantum eraser effect [11] or complementarity principle [12], by noting the inherent relation between entanglement and quantum interference. It is found that, by properly designing the quantum eraser composed by the entangled electron pair, the concurrence, i.e. degree of entanglement, can be obtained by the amplitude of Aharonov-Bohm (AB) oscillation of the current correlation [10].

In the original proposal of the quantum eraser method, stationary AB effect and pure entangled states are considered [10]. In this paper, we generalize our exploration to the nonstationary AB effect and mixed states caused by spin dephasing. In particular, we focus on a novel quantum spin Hall system (QSH) [13], for the helical property of edge states in QSH makes the proposal more




FIG. 1. Illustration of the proposed setup. The entangled electron pairs are tunneling separately from the entangler (labeled by E) into the helical edge channels of QSHs. The spin up (down) channels are sketched as solid (dashed) lines. The narrow regions in both sides bring coupling between the upper and lower edge channels, and serve as beam splitters. In the right side, a magnetic flux Φ is enclosed by the loop formed by the upper and lower channels. Four terminals ($L_{1,2}, R_{1,2}$) measure the electronic currents and current correlations.

realizable. Firstly, the spin and moving direction of helical electrons are bounded, which leads to an all-electrical-control of electron spin [8]. Secondly, the back scattering is forbidden by the time reversal symmetry, so the spin relaxation time is extremely long.

The proposed experimental setup is shown in Fig. 1, where the entangler in the middle region is weakly coupled to two QSHs in both sides. When a voltage bias is applied between the entangler and terminals, entangled electron pairs will tunnel into the helical edge channels of QSHs. They may be injected into the same QSH, or they may be separately injected into different QSHs. However, we here only consider the latter process, because the injection into the same QSH has only a higher order contribution to the current correlation in the weak coupling limit [8, 14]. In each QSH, there is a narrow region with the opposite edge channels being coupled, which leads to the scattering between edges. It has been demonstrated that by tuning the gate voltage properly, the quantum

point contact structure can serve as an ideal beam splitter without any back scattering [15], which is beneficial to entanglement detection. We here consider that a time-dependant flux $\Phi(t)$ is applied in the right QSH of Fig. 1, which leads to the nonstationary AB effect.

The entangler can be realized by using superconductors [5, 6] or quantum dots [16]. Here we adopt the former as an example, while the theory itself is actually general, which can be extended to other kinds of entanglers. We assume that the superconductor is grounded and a bias voltage $-eV$ ($<$ the superconducting gap Δ) is applied to all the terminals $L_{1,2}, R_{1,2}$, so the Cooper pairs are split into edge channels of the QSHs [17] with a tunneling coefficient $\chi \ll 1$. The entangled states may be expressed as [8, 14]

$$|\Psi\rangle = \chi \int_{-eV}^{eV} d\varepsilon \left[\sqrt{\kappa} a_{L\uparrow}^\dagger(\varepsilon) a_{R\downarrow}^\dagger(-\varepsilon) \mp \sqrt{1-\kappa} a_{L\downarrow}^\dagger(\varepsilon) a_{R\uparrow}^\dagger(-\varepsilon) \right] |0\rangle, \quad (1)$$

where $|0\rangle$ is the zero temperature ground state of filled Fermi sea with a Fermi energy $-eV$, and $a_{i,\sigma}^\dagger(\varepsilon)$ creates an electron with energy ε and spin σ in the i side which satisfies the anticommutation relation $\{a_{i,\sigma}(\varepsilon), a_{j,\sigma'}^\dagger(\varepsilon')\} = \delta_{i,j} \delta_{\sigma,\sigma'} \delta(\varepsilon - \varepsilon')$. The helicity of the edge states indicates the spin-edge correspondence. For example, $a_{L\uparrow}$ describes an electron moving left along the lower edge with its spin polarized up (Fig. 1). Though we here focus on the singlet and triplet entangled states, we keep the parameter κ tunable to show clearly how the physical results changes continuously from the mostly entangled states ($\kappa = 1/2$) to product states ($\kappa = 0, 1$). For the pure state in Eq. (1), the degree of entanglement can be described by the concurrence $\mathcal{C} = 2\sqrt{\kappa(1-\kappa)}$ [18]. Specifically, when $\mathcal{C} = 1$, the “ $\mp$ ” in Eq. (1) correspond to the singlet and triplet entangled states, respectively.

After transmission into QSHs, the electrons get scattered at the narrow regions as in Fig. 1, before they finally arrive at the terminals. They may change their edges during the scattering processes, while the back scattering is completely ruled out by tuning the gate voltages properly [15]. Thus the electron operators in L_1, R_1 can be written as

$$\begin{aligned} a_{L1} &= t_{2L} a_{L\uparrow} + t_{1L} a_{L\downarrow}, \\ a_{R1} &= t_{1R} a_{R\uparrow} + t_{2R} a_{R\downarrow}, \end{aligned} \quad (2)$$

where the coefficients $t_{1,2i}$ describe the amplitudes for the same-edge and cross-edge transmissions in the i side, respectively. The electrons in the other two terminals L_2, R_2 can be described as well, which are omitted here.

In order to probe entanglement, we calculate the zero frequency current correlation between terminals L_1, R_1 , which can be expressed by the noise power

$$S = 2 \int_{-\infty}^{\infty} dt [\langle \hat{I}_{L1}(t) \hat{I}_{R1}(0) \rangle - \langle \hat{I}_{L1}(t) \rangle \langle \hat{I}_{R1}(0) \rangle], \quad (3)$$

where the current operator defined as $\hat{I}_{i1}(t) = (e/h) \times \int \int dE dE' e^{i(E'-E)t/\hbar} a_{i1}^\dagger(E') a_{i1}(E)$ and the higher order term $\langle \hat{I}_{L1}(t) \rangle \langle \hat{I}_{R1}(0) \rangle \sim \chi^4$ can be neglected. By utilizing Eq. (2) and the anticommutation relation of Fermion operators, we can obtain the time dependant current correlation after some algebra

$$\begin{aligned} \langle \hat{I}_{L1}(t) \hat{I}_{R1}(0) \rangle &= \left(\frac{e\chi}{h}\right)^2 \int_{-eV}^{eV} d\varepsilon \int_{-eV}^{eV} d\varepsilon' e^{i(\varepsilon-\varepsilon')t/\hbar} \times \\ &\quad \left[\kappa T_{2L} T_{2R} + (1-\kappa) T_{1L} T_{1R} \mp \mathcal{C} \sqrt{T_{2L} T_{2R} T_{1L} T_{1R}} \cos \varphi(t) \right], \end{aligned} \quad (4)$$

where the transmission probabilities are $T_{1,2i} = |t_{1,2i}|^2$ and the total phase is defined as $\varphi(t) = \text{Arg}[t_{2L}^* t_{1L} t_{2R}^* t_{1R}]$. From Eq. (4), we notice that the current correlation can be tuned by the flux in the right QSH (Fig. 1), and the strongest interference occurs when $T_{1,2L} = T_{1,2R} = 1/2$, which will be adopted in the following.

In order to investigate both *dc* and *ac* AB effects, a harmonic flux $\Phi(t) = \Phi_a \sin \Omega t$ with amplitude Φ_a is introduced [19], so the phase has the form $\varphi(t) = \varphi_0 + 2\pi(\Phi_a/\Phi_0) \sin \Omega t$ with a constant term φ_0 including all stationary phase accumulated by transport, scattering and also the *dc* component of the flux, where $\Phi_0 = hc/e$ is the flux unit.

Given the periodicity of the phase, $\varphi(t+2\pi/\Omega) = \varphi(t)$, the interference term can be expanded as Fourier series by utilizing the identity

$$e^{i2\pi \frac{\Phi_a}{\Phi_0} \sin \Omega t} = \sum_{n=-\infty}^{\infty} J_n(2\pi \frac{\Phi_a}{\Phi_0}) e^{in\Omega t}, \quad (5)$$

where J_n is the Bessel function of the first kind. Inserting Eq. (5) into Eqs. (3, 4), and performing all the integrals over t, ε and ε' , we finally obtain an elegant expression for the differential noise (DN) $\Lambda = \partial S / \partial eV$ as

$$\Lambda = \chi^2 \frac{e^2}{h} \left\{ 1 \mp \mathcal{C} \cos \varphi_0 \left[J_0 + 2 \sum_{n>0} J_{2n} \theta(eV - n\hbar\Omega) \right] \right\}, \quad (6)$$

where we have omitted the variable $2\pi \frac{\Phi_a}{\Phi_0}$ of the Bessel functions for simplicity and θ is the Heaviside stepfunction. Eq. (6) is a main result of the present work, which contains important information for probing the concurrence.

DC case. First we consider the case $\Phi_a = 0$ when the *ac* component of the flux is absent. In this case, we have $J_0 = 1, J_{2n} = 0$ in Eq. (6), so the expression is just an ordinary AB oscillation of the DN as $\Lambda = \chi^2 e^2 / h (1 \mp \mathcal{C} \cos \varphi_0)$ [10], which is sketched by Fig. 2 for the triplet entangled states. The relation between entanglement and current correlation is reflected by the equality of the visibility defined as $\nu = (\Lambda_{\max} - \Lambda_{\min}) / (\Lambda_{\max} + \Lambda_{\min})$ and

the concurrence, namely $\nu = \mathcal{C}$. The mechanism of this equality is the quantum eraser effect [11]. Noting that the original state by Eq. (1) has the typical form expressed simply as $\sqrt{\kappa} |\uparrow\downarrow\rangle \mp \sqrt{1-\kappa} |\downarrow\uparrow\rangle$, where the right electron transports in the interference loop and the left electron is the entangled partner. There are only two transport channels contributing to the interference without any spin degeneracy, as a result of the helical property of electrons in QSH. Now if only the right electron is measured, there is no AB oscillation of current in R_1 . The reason is that the left electron serves as the which way marker of the right one. For example, if the left electron is in the state $|\uparrow\rangle$, then we can infer that the right electron picks the lower channel. According to the complementarity principle [12], whenever the which way information is marked, the interference will not occur. However, if we make a correlated measurement between the two electrons, or equivalently, the current correlation between L_1, R_1 , then the result will be distinctly different. The fantastic feature of entanglement lies in that as long as the spin of the left electron gets measured, the right electron will collapse into a corresponding state, which may contain no path information at all and the interference will recur. To understand this point clearly, we can take $t_{1,2L} = 1/\sqrt{2}$ as an example, and the spin state detected by L_1 terminal is $|L_1\rangle = 1/\sqrt{2}(|\uparrow\rangle + |\downarrow\rangle)$, with the original spin information $|\uparrow\rangle, |\downarrow\rangle$ totally erased. Now if an electron is detected by L_1 , we can infer from the original entangled state Eq. (1) that the amplitude of its entangled partner detected by R_1 is $t_{2R}\sqrt{\kappa} \mp t_{1R}\sqrt{1-\kappa}$, where the interference recurs. It is also clear that the amplitude of AB oscillation gives out exactly the degree of entanglement, because the quantum eraser effect is a direct result of nonlocal quantum correlation from entanglement [10].

AC case. Then we set $\varphi_0 = 0$ to investigate the non-stationary AB effect, and the DN has the form $\Lambda = \chi^2 e^2/h \{1 \mp \mathcal{C} [J_0 + 2 \sum_{n>0} J_{2n} \theta(eV - n\hbar\Omega)]\}$. The main characteristic of the result is the step structures as shown in Fig. 3. As long as the voltage bias V increases by $\hbar\Omega/e$, the DN jumps onto a new step. The increase (decrease) of the n th step is $2\mathcal{C}J_{2n}$ ($n > 0$) for triplet (singlet) state. Remarkably, through the measured ratios of heights of these steps, one can first obtain the flux amplitude Φ_a . Then with the fixed Φ_a , the concurrence can be measured directly through the heights of these steps. In Fig. 3(a), we present the step structures of Λ for triplet entangled states ($\mathcal{C} = 1$), when the flux amplitude Φ_a takes different values. The figures of the singlet states are simply the mirror symmetry about $\chi^2 e^2/h$, which comes from the “ $\mp$ ” in the original states in Eq. (1), indicating that they are distinguishable by the step shapes. We also note that when the voltage bias is big enough, the height of the steps will approach 0 and $2\chi^2 e^2/h$ for singlet and triplet states, respectively. This is not surprising if we notice the identity $\sum_{n=-\infty}^{\infty} J_n = 1$.




FIG. 2. Plot of the DN as a function of stationary flux φ_0 for triplet entangled states (The singlet states are the mirror symmetry about $\chi^2 e^2/h$). The solid lines stand for pure states with the dephasing parameter $\gamma = 1$ and the dashed and dotted lines represent the mixed states with $\gamma = 0.5, 0.3$.




FIG. 3. Plot of the DN as a function of voltage bias for the triplet entangled states (The singlet states are also the mirror symmetry about $\chi^2 e^2/h$). (a) The pure triplet entangled states with the amplitudes of the ac fluxes $\Phi_a = 0.5\Phi_0, \Phi_0, 1.5\Phi_0$ corresponding to the lines 1, 2, 3, respectively. (b) For a fixed amplitude of flux $\Phi_a = \Phi_0$, with different dephasing parameter $\gamma = 1, 0.5, 0.3$.

Spin dephasing It is worthwhile to discuss the spin dephasing effect in the QSHs. The spin flipped back scattering is totally suppressed in the helical edge channels by the time reversal symmetry, so theoretically, the spin relaxation process can be neglected in the present model with nano-scale. On the other hand, the spin dephasing process always exists, and due to the helical property, the spin and edge freedoms are bounded together. Therefore any interaction between edge and environment that introduces a phase uncertainty on the edge freedom will also lead to the spin dephasing. Here we would like to

give a general discussion on the spin dephasing effect and its effect on the AB effect of current correlation, without limiting in a specific environment, such as a free electromagnetic field and the charge fluctuation in QSH, etc.

When the electron interacts with the environment, they together are generally in an entangled state. The interference of the electron can be obtained by performing integral over all the freedoms of the environment. In the quasi-classical approximation, one finds that the effect of the interaction is only to multiply the interference term in Eq. (4) by a scalar product of the initial and final states of the environment as [20] $\gamma = \langle \xi_0 | \hat{T} \exp[-i/\hbar \int_0^{\tau_0} dt V_I(t)] | \xi_0 \rangle$, during the time τ_0 of electron traversing through the system, where $|\xi_0\rangle$ is the initial state of the environment and $V_I(t)$ is the interaction term in the interaction picture. Usually the phase accumulation in many-body environment satisfies Markov approximation, i.e. $[V_I(t), V_I(t')] = 0$, for a short memory time of environment compared with τ_0 . By using the central limit theorem, the dephasing parameter is an exponential decay with the phase uncertainty as $\gamma(\tau_0) = e^{-1/2 \langle \delta\varphi^2 \rangle}$ [20]. Given that the nonstationary case discussed here, it is necessary to compare two time scales, i.e. the electron traversing time τ_0 and the period of the *ac* flux $2\pi/\Omega$. Only when $\Omega\tau_0 \ll 2\pi$ is satisfied, the phase $\varphi(t)$ can be treated quasi-classically as a function of time in Eq. (4). Taking the HgTe quantum wells as an example [13], the Dirac velocity of edge states is given by $v \simeq 5.5 \times 10^5$ m/s, so the electron spends about $\tau_0 = l/v \simeq 1.8$ ps to transport through the QSH for a nano-scale system with a typical length $l \sim 1\mu\text{m}$. As a result, the quasi-classical condition is $\Omega \ll 3.5\text{THz}$, which is fulfilled by the usual microwave frequency $\Omega \sim 1\text{GHz}$ or by the time-dependant gate voltage on the edge channel [19], which results in the *ac* AB effect.

Based on the above discussions, the dephasing effect can be effectively described by the dephasing parameter γ within the interval $0 \leq \gamma < 1$, which changes the pure states into mixed states. For the singlet and triplet entangled states in Eq. (1), the original pure states $1/\sqrt{2}(|\uparrow\downarrow\rangle \mp |\downarrow\uparrow\rangle)$ is now substituted by the density matrix $\hat{\rho} = 1/2[|\uparrow\downarrow\rangle\langle\uparrow\downarrow| + |\downarrow\uparrow\rangle\langle\downarrow\uparrow| \mp \gamma(|\uparrow\downarrow\rangle\langle\downarrow\uparrow| + |\downarrow\uparrow\rangle\langle\uparrow\downarrow|)]$, with the concurrence $\mathcal{C} = \gamma$ [21]. The current correlation is now calculated through the formula $\langle \hat{I}(t)\hat{I}(0) \rangle = \chi^2 \int_{-eV}^{eV} d\varepsilon \int_{-eV}^{eV} d\varepsilon' \text{Tr}[\hat{\rho}\hat{I}(t)\hat{I}(0)]$, and finally we can obtain exactly the same equation Eq. (6), merely with the concurrence $\mathcal{C} = \gamma$. Therefor, we conclude that no matter the spin dephasing exists or not, the DN formula Eq. (6) is generally applicable, with $\mathcal{C} = 1, \gamma$ corresponding to pure and mixed states. The numerical results can be found in Fig. 2 and Fig. 3(b) for *dc* and *ac* cases, respectively. In the *dc* case, the visibility ν is suppressed by the spin dephasing effect, while the equality $\nu = \mathcal{C}$ between visibility and concurrence is general, for both pure and mixed states. In the *ac* case, the height of

each step is also suppressed by the dephasing parameter γ , while all the arguments for pure states are still valid, only with $\mathcal{C} = 1$ being replaced by $\mathcal{C} = \gamma$. Therefor, the dephasing effect in QSHs can also be measured through the *dc* and *ac* AB effect of current correlation.

In summary, we have studied both *dc* and *ac* AB effects of the current correlation contributed by entangled electron pairs in QSHs. It has been shown that for singlet and triplet entangled states, the degree of entanglement can be probed by the visibility of the stationary AB oscillation, or the step structures in the nonstationary case, no matter the spin dephasing exists or not, providing a new method for entanglement detection.

One of the authors (W. C.) would like to thank his friend Wei Xia for his tremendous support and encouragement. This work was supported by 973 Program (Grants No. 2011CB922100, No. 2011CBA00205, and No. 2009CB929504), by NSFC (Grants No. 11074111, No. 11174125, and No. 11023002), by PAPD of Jiangsu Higher Education Institutions, by NCET, by the Fundamental Research Funds for the Central Universities, by the GRF (HKU7058/11P), CRF (HKU-8/11G) of the RGC of Hong Kong, and the URC fund of HKU.

* zwang@hku.hk

† shen@nju.edu.cn

- [1] A. Einstein, B. Podolsky, and N. Rosen, *Phys. Rev.* **47**, 777 (1935); E. Schrödinger, *Proc. Cambridge Philos. Soc.* **31**, 555 (1935).
- [2] J. S. Bell, *Physics* (Long Island City, N.Y.) **1**, 195 (1964); J. S. Bell, *Rev. Mod. Phys.* **38**, 447 (1966).
- [3] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, England, 2000).
- [4] D. Loss and D. P. DiVincenzo, *Phys. Rev. A* **57**, 120 (1998).
- [5] L. Hofstetter, S. Csonka, J. Nygård, and C. Schönenberger, *Nature* (London) **461**, 960 (2009); L. G. Herrmann, F. Portier, P. Roche, A. L. Yeyati, T. Kontos, and C. Strunk, *Phys. Rev. Lett.* **104**, 026801 (2010); J. Wei and V. Chandrasekhar, *Nature Phys.* **6**, 494 (2010).
- [6] G. B. Lesovik, T. Martin, and G. Blatter, *Eur. Phys. J. B* **24**, 287 (2001); P. Recher, E. V. Sukhorukov, and D. Loss, *Phys. Rev. B* **63**, 165314 (2001).
- [7] S. Kawabata, *J. Phys. Soc. Jpn.* **70**, 1210 (2001); N. M. Chtchelkatchev, G. Blatter, G. B. Lesovik, and T. Martin, *Phys. Rev. B* **66**, 161320(R) (2002).
- [8] W. Chen, R. Shen, L. Sheng, B. G. Wang, and D. Y. Xing, *Phys. Rev. Lett.* **109**, 036802 (2012).
- [9] G. Burkard, D. Loss, E. V. Sukhorukov, *Phys. Rev. B* **61**, R16303 (2000).
- [10] W. Chen, R. Shen, Z. D. Wang, L. Sheng, B. G. Wang and D. Y. Xing, *Phys. Rev. B* **87**, 155308 (2013).
- [11] M. O. Scully and K. Drühl, *Phys. Rev. A* **25**, 2208 (1982).
- [12] T. J. Herzog, P. G. Kwiat, H. Weinfurter, and A. Zeilinger, *Phys. Rev. Lett.* **75**, 3034 (1995).
- [13] M. Z. Hasan and C. L. Kane, *Rev. Mod. Phys.* **82**, 3045

- (2010); X. L. Qi and S. C. Zhang, *Rev. Mod. Phys.* **83**, 1057 (2011).
- [14] P. Samuelsson, E. V. Sukhorukov, and M. Büttiker, *Phys. Rev. Lett.* **91**, 157002 (2003).
 - [15] V. Krueckl and K. Richter, *Phys. Rev. Lett.* **107**, 086803 (2011).
 - [16] D. Loss and E. V. Sukhorukov, *Phys. Rev. Lett.* **84**, 1035 (2000); W. D. Oliver, F. Yamaguchi, and Y. Yamamoto, *Phys. Rev. Lett.* **88**, 037901 (2002).
 - [17] K. Sato, D. Loss, and Y. Tserkovnyak, *Phys. Rev. Lett.* **105**, 226401 (2010); W. Chen, R. Shen, L. Sheng, B. G. Wang, and D. Y. Xing, *Phys. Rev. B* **84**, 115420 (2011).
 - [18] S. Hill and W. K. Wootters, *Phys. Rev. Lett.* **78**, 5022 (1997); W. K. Wootters, *Phys. Rev. Lett.* **80**, 2245 (1998).
 - [19] The flux $\Phi(t)$ can also be replaced by a gate voltage $V_g(t)$ on one edge channel of the interference loop, through the relation $\Phi(t) = V_g(t)d\Phi_0/hv$, with d the length of the gating region and v the Dirac velocity. Due to the helicity of the edge states, the gate voltage will not lead to any back scattering, and only contributes an extra phase, as the same role played by the flux.
 - [20] A. Stern, Y. Aharonov, Y. Imry, *Phys. Rev. A* **41**, 3436 (1990).
 - [21] The concurrence of mixed states can be calculated due to Wootters [18] by the formula $\mathcal{C} = \max\{0, \sqrt{\lambda_1} - \sqrt{\lambda_2} - \sqrt{\lambda_3} - \sqrt{\lambda_4}\}$, where $\lambda_1 \geq \lambda_2 \geq \lambda_3 \geq \lambda_4$ are the eigenvalues of the matrix product $\rho(\sigma_y \otimes \sigma_y)\rho^*(\sigma_y \otimes \sigma_y)$.